

Primefact

Hydatids – The Basics

March 2026, PUB17/553[v2], Third Edition
Animal Biosecurity, NSW DPIRD

Introduction

In Australia, hydatid disease is caused by the hydatid tapeworm, which infects dogs, dingoes and foxes. Humans can also be intermediate hosts, with serious health consequences.

Losses from hydatid disease in livestock arise from the downgrading of edible meat by-products because of hydatid cysts.

Control is by preventing or eliminating hydatid tapeworm infections in the final hosts – dogs, foxes and dingoes. Controlling hydatids will also help to control sheep measles, caused by a tapeworm with a similar lifecycle.

Hydatid Lifecycle

The *E. granulosus* tapeworm consists of 3-4 segments and is up to 6 mm long. Thousands of these small tapeworms can inhabit the intestines

of a dog without ill effect. Tapeworm segments containing eggs are periodically shed in the faeces. The eggs, which are resistant to weathering, are scattered by wind and water.

Susceptible intermediate hosts swallow the eggs. Such hosts include sheep, pigs, goats, camels, deer, cattle, horses, kangaroos and wallabies, and humans.

Swallowed eggs – for example, from pasture or, for humans, unwashed hands or vegetables – yield embryos which traverse the gut wall and are transported to various tissues, usually the liver

or lungs, but sometimes the brain. Hydatid cysts (fluid filled sacs) then develop. The next generation of tapeworm heads develop within these cysts.

The cycle is completed when final hosts (dogs, dingoes, foxes) consume tissues with hydatid cysts or contaminated with cyst fluid.

The sylvatic (wildlife) life cycle may involve dingoes, dogs and foxes as final hosts and kangaroos and wallabies as intermediate hosts.

Controlling hydatids

- Worm your dogs (working dogs and pets) monthly with a de-worming product that states on the label it is effective against Hydatids.
- Never feed raw offal (liver/lungs/hearts/kidneys) from sheep, goats, cattle or any species of wildlife to dogs. Freeze it for at least 14 days at -10° to -15°C or cook it thoroughly.
- Secure dogs at night and when not working so they do not wander and scavenge on carcasses.
- Put all offal in a dog-proof pit, and where possible dispose of the bodies of stock that have died in the paddock.

Engage in good personal hygiene; always ensure you and your family wash your hands after handling dogs and especially before you eat.

Eliminate infections

The only suitable tape wormer for treating dogs is praziquantel. This active ingredient is found in many dog wormers (check the label).

When treating potentially infected dogs, take great care in disposing (by deep burial, or burning) of droppings for three days after treatment.

Dogs at on-going risk of infection will need to be treated with praziquantel every 6 weeks.

Hygiene

As with many zoonoses (diseases transmissible between animals and humans), attention to hygiene can markedly reduce the risk of infection.

Always thoroughly wash your hands before putting your hands near your mouth (for eating, cigarette smoking, etc.), especially after handling dogs or being in potentially contaminated areas such as around kennels. Wash vegetables and fruit before eating or cooking them. Teach your children to do the same, by instruction and by example.

More information

- Tel: 1800 680 244, Animal Biosecurity and Welfare
- Department of Primary Industries and Regional Development at animal.biosecurity@dpir.nsw.gov.au,
- Local Land Services, Tel: 1300 795 299
- To view details of your LLS region: <http://www.lls.nsw.gov.au/>

© State of New South Wales through DPIRD March 2026. You may copy, distribute, display, download and otherwise freely deal with this publication for any purpose, provided that you attribute the Department of Primary Industries and Regional Development as the owner. However, you must obtain permission if you wish to charge others for access to the publication (other than at cost); include the publication in advertising or a product for sale; modify the publication; or republish the publication on a website. You may freely link to the publication on a departmental website.

ISSN: 1832-6668

Disclaimer: The information contained in this publication is based on knowledge and understanding at the time of writing (March 2026) and may not be accurate, current or complete. The State of New South Wales (including the DPIRD), the author and the publisher take no responsibility, and will accept no liability, for the accuracy, currency, reliability or correctness of any information included in the document (including material provided by third parties). Readers should make their own inquiries and rely on their own advice when making decisions related to material contained in this publication.